


Pazopanib (Votrient®) Medication

Pazopanib (the generic name for Votrient) is a cancer medicine you take by mouth.

How to Take This Medicine

You take four pazopanib pills once a day without food. Each pill is 200 milligrams. The total dose will be 800 milligrams once a day, at the same time each day.



It is very important that you take this medicine on an empty stomach with water only.

Do not eat for at least 2 hours *before* taking the medicine and for at least 1 hour *after* taking it.

EXAMPLES

If you take your dose each morning:

Take four pills when you wake up first thing in the morning, when your stomach is empty. Wait one hour before eating breakfast or drinking anything but water.

If you take your dose each night:

Wait at least two hours after eating dinner or any dessert. Take four pills with water before you go to bed, then do not eat again until breakfast the next morning.

Important Warnings

Some foods and other medicines could interact with pazopanib or keep it from working. These include common antacids and drugs to lower cholesterol. If another doctor prescribes you medication for any reason, make sure he or she knows you are taking pazopanib.

- DO NOT take St. John's Wort.
- DO NOT eat or drink grapefruit or grapefruit juice.
- CHECK with your doctor before taking antacids or any other medicine while taking pazopanib.

Common Side Effects

These are the most common side effects of pazopanib:

- Diarrhea
- High blood pressure - Have your doctor check your blood pressure regularly
- Hair color changes - Hair may lose its color during treatment
- Decreased thyroid levels - This will be monitored with blood tests
- Nausea
- Fatigue
- Mucositis - This may include a sensitive mouth, sores in the mouth, or changes in taste
- Hand-foot syndrome - This includes skin changes such as redness, blisters, calluses, cracking, flaking, pain, or sensitivity on your palms or feet

When to Call Your Doctor

For your safety, please call our office to be sure it is OK to take any other medication while taking pazopanib. This includes any kind of antacid.

If you have any of the side effects listed above, please call us. We can give you medicines to help manage side effects. Your doctor will also ask about side effects during each visit.

Tell us if you are scheduled to have any type of procedure or surgery. You may need to stop taking pazopanib for a short time.

Monday–Friday, 8 a.m.–5 p.m.

Call your doctor's office at this number:

**After hours, weekends, and holidays, call
801-587-7000.**

Ask for the oncologist on call.